

Energy dependence of string fragmentation function and ϕ meson production

Ben-Hao Sa^{1,2,4,5*}, Xu Cai^{4,2}, Chinorat Kobdaj³,
Zhong-Qi Wang¹, Yu-Peng Yan³, and Dai-Mei Zhou²

¹ *China Institute of Atomic Energy,*

P. O. Box 275 (18), Beijing, 102413 China

² *Institute of Particle Physics, Huazhong Normal University, Wuhan, 430079 China*

³ *School of Physics, Suranaree University of Technology,
Nakhon Ratchasima 30000, Thailand*

⁴ *CCAST (World Lab.), P. O. Box 8730 Beijing, 100080 China*

⁵ *Institute of Theoretical Physics, Academia Sinica, Beijing, 100080 China*

Abstract

The ϕ meson productions in $Au + Au$ and/or $Pb + Pb$ collisions at AGS, SPS, RHIC, and LHC energies have been studied systematically with a hadron and string cascade model LUCIAE. After considering the energy dependence of the model parameter α in string fragmentation function and adjusting it to the experimental data of charged multiplicity to a certain extent, the model predictions for ϕ meson yield, rapidity, and/or transverse mass distributions are compatible with the experimental data at AGS, SPS and RHIC energies. A calculation for $Pb + Pb$ collisions at LHC energy is given as well. The obtained fractional variable in string fragmentation function shows a saturation in energy dependence. It is discussed that the saturation of fractional variable in string fragmentation function might be a qualitative representation of the energy dependence of nuclear transparency.

PACS numbers: 25.75.Dw, 24.10.Lx, 24.85.+p, 25.75.Gz

* E-mail: sabh@iris.ciae.ac.cn

Strangeness enhancement was suggested in the early eighties [1] as one of the most promising signatures for the creation of a Quark-Gluon Plasma (QGP) phase in relativistic nuclear collisions. Following the experimental observations on strangeness enhancement in proton-nucleus and nucleus-nucleus collisions at the SPS energies [2, 3, 4, 5, 6, 7] the WA97 has measured a clear enhancement of multi-strange baryons (Λ, Ξ, Ω) with their strange quark content in 158A GeV/c $Pb + Pb$ collisions relative to $p + Pb$ collisions [8]. Recently the STAR data on the strangeness production in $Au + Au$ collisions at $\sqrt{s_{nn}}=130$ GeV were reported [9].

TABLE I: Global hadron multiplicity in relativistic nucleus-nucleus collisions

Reaction		$Au + Au$	$Pb + Pb$	$Au + Au$	$Pb + Pb$
$\sqrt{s_{nn}}$ (GeV)		4.66	17.3	130	5500
Centrality	Exp.	$\leq 4\%$	$\leq 4\%$	$\leq 5\%$ (BRAHMS)	$\leq 10\%$
	LUCIAE	$b \leq 2.82$ fm	$b \leq 3.5$ fm	$b \leq 3.2$ fm	$b \leq 4.54$ fm
N_{ch}	Exp.			$3860 \pm 300^{1)}$	
	LUCIAE			4114	$15646^{2)}$
$(\langle \pi^+ \rangle + \langle \pi^- \rangle)/2$	Exp.	$105^{3)}$	$611^{4)}$		
	LUCIAE	102	611		$5851^{2)}$
α		0.4	1.3	12	18
$\langle z \rangle$		0.286	0.565	0.923	0.947

1. $-4.7 < \eta < 4.7$, from [29]

2. in full phase space

3. $0.6 < y < 2.6$, taken from [30]

4. cf. [14]

As the mesonic counterpart, the enhancement of ϕ meson production in relativistic nuclear collisions was also suggested as an evidence of the QGP formation in Ref. [10], since in the environment of a QGP the copious strange and antistrange quarks originating from gluon annihilation would be very likely to coalesce forming ϕ mesons during the hadronization period. Due to the small cross sections of ϕ mesons interacting with non-strange hadrons [1, 10], penetrating ϕ mesons are also messengers of the early stage of the colliding system. Thus, the ϕ meson is not only a promising signature for the QGP formation but also a good probe to study the reaction dynamics. After the experimental observations on ϕ meson productions in $Au + Au$ collisions at AGS [11] and sulfur-nucleus collisions and $Pb + Pb$ collisions at SPS [12, 13, 14], the STAR collaboration




FIG. 1: Rapidity (left panel) and transverse mass (right panel) distributions of ϕ mesons in $Pb+Pb$ collisions at 158A GeV/c

reported recently the

at $\sqrt{s_{nn}}=130 \text{ GeV}$ [15].




FIG. 2: Transverse mass distributions ($-0.5 < y < 0.5$) of ϕ mesons in $Au+Au$ collisions at $\sqrt{s_{nn}}=130 \text{ GeV}$

The model studies on the ϕ meson enhancement in relativistic nucleus-nucleus collisions are rare [16, 17, 18, 19] and there exists up to now no theoretical investigation on the energy dependence of ϕ production from AGS, to SPS, to RHIC, and up to LHC energies systematically. In this letter, a

hadron and string cascade model, LUCIAE [20] was employed to investigate first time the energy dependence of ϕ meson production systematically. We have successfully used LUCIAE to study the enhanced production of multi-strange baryons (Λ, Ξ, Ω) and determined the model parameters related to the production of strange particles [21, 22].

TABLE II: ϕ mesons yield in relativistic nucleus-nucleus collisions

Reaction	$\sqrt{(s_{nn})}$ (GeV)	Centrality		$\langle N_\phi \rangle$	
		Exp.	LUCIAE	Exp. data	LUCIAE
$Au + Au$	4.66	301 ¹⁾	307 ¹⁾	0.252 $\pm$ 0.107 ²⁾	0.14
$Pb + Pb$	17.3	$\leq 4\%$	$b \leq 3.5$ fm ³⁾	7.6 $\pm$ 1.1 ³⁾	6.48
$Au + Au$	130	$\leq 11\%$ (STAR)	$b \leq 4.31$ fm	5.73 $\pm$ 1.06 ⁴⁾	5.10
$Pb + Pb$	5500	$\leq 10\%$	$b \leq 4.54$ fm		132 ⁵⁾

1. $\langle N_{part} \rangle$
2. $\frac{dN_\phi}{dy}$ within $0.9 < y < 1.4$ [11]
3. cf. [14]
4. $\frac{dN_\phi}{dy}$ within $-0.5 < y < 0.5$ [15]
5. full phase space

The LUCIAE model is based on FRITIOF [23], which is an incoherent hadron multiple scattering and string fragmentation model. In FRITIOF, the nucleus-nucleus collision is depicted simply as a superposition of nucleon-nucleon collisions. What characterizes LUCIAE beyond FRITIOF are the following features: First of all, the rescattering among the participant and spectator nucleons and the produced particles from string fragmentation are generally taken into account [24]. However, as proposed in [1, 10] the effects of the final state interactions on the ϕ meson production and propagation are neglected. Secondly, the collective effect in the gluon emission of strings is considered by firecracker model [25]. Thirdly, a phenomenological mechanism for the reduction of the s quark suppression in the string fragmentation is introduced [21] resulting the effective string tension and therefore the pertained JETSET parameters vary automatically with the energy, centrality, and size of reaction system. Those JETSET parameters: parj(2), (3), (1), and (21) in program are, respectively, the suppression of s quark pair production in string fragmentation compared with u or d pair production, the extra suppression of strange diquark production compared with the normal suppression of strange quarks, the suppression of diquark-antidiquark pair production compared with quark-antiquark pair production, and the width in the Gaussian




FIG. 3: (a) fractional variable in string fragmentation function $\langle z \rangle$ and (b) $\langle N_{CH} \rangle / \sqrt{s_{AA}}$ as a function of $\sqrt{s_{nn}}$

From the AGS energy ($\sqrt{s_{nn}} \sim 5$ GeV) to SPS ($\sqrt{s_{nn}} \sim 20$ GeV) to RHIC ($\sqrt{s_{nn}} \sim 200$ GeV) and then to the LHC energy ($\sqrt{s_{nn}} \sim 5500$ GeV) there are four energy magnitudes spanned. Among the nucleus-nucleus collisions in above energy region their dynamic behaviors, of course, must be different observably from each other. To describe the nucleus-nucleus collisions in such a wide energy region one either selects suitable model for different energy or considers the energy dependence of model ingredients properly if one single model is used. This idea has indeed been adopted in e^+e^- studies quite early [27]. Recently the thermal model with two main parameters (temperature T and baryon chemical potential μ_B) was used to fit the experimental data of global hadronic yields for nucleus-nucleus collisions at SIS ($\sqrt{s_{nn}} \sim 1$ GeV), AGS, SPS, and RHIC energies, the obtained values of T and μ_B are quite different from each other [28]. Therefore, for dynamical models it is also needed to consider the energy dependence of the involved dynamical ingredients properly in order to describe well the nucleus-nucleus collisions in region spanned four energy magnitudes. Of course, some dynamical ingredients might be sensitive to the energy and others might not be. Many dynamical models (including LUCIAE) working well for nucleus-nucleus collisions at SPS energy, however, overestimate the charged multiplicity at RHIC energy. One reason might just be that there are ingredients either energy independent or energy dependent without taken into account properly.

In LUCIAE model (in FRITIOF and in JETSET eventually) a very important dynamical ingredient, the string fragmentation function, is energy independent. Thus the LUCIAE model is

extended in this letter to take the energy dependence of string fragmentation function into account. As mentioned in [26] that the string fragmentation function $f(z)$, which expresses the probability that a given fractional variable z is sampled, could be arbitrary in principle. In [26] several such choices are given, besides the one of Lund fragmentation function. For describing easily the relativistic nucleus-nucleus collisions over four energy magnitudes one of the simplest option for the string fragmentation function in [26] is employed here for the moment. That string fragmentation function reads

$$f(z) = \alpha(\sqrt{s_{nn}}) \times z^{\alpha(\sqrt{s_{nn}})-1}, \quad (1)$$

where the fractional variable z refers to the fraction of light-cone momentum taken away by produced hadron from the fragmenting string, the parameter α in this string fragmentation function is assumed to be energy dependent now.

We first ran LUCIAE for $Pb+Pb$ and $Au+Au$ collisions, respectively, at SPS and RHIC energies. In LUCIAE calculations the centrality cut and the windows of rapidity (pseudorapidity) and p_t were the same as those in corresponding experiments (the same later). Adjusting the parameter $\alpha(\sqrt{s_{nn}})$ in order that the global hadron multiplicity from LUCIAE could be compatible with the corresponding experimental data, the obtained LUCIAE results are given in Tab. 1 together with the experimental data. The data of $Pb + Pb$ collisions at SPS energy were taken from [14] and $Au + Au$ collisions at $\sqrt{s_{nn}}=130$ GeV from [29]. Since in LUCIAE the reduction mechanism of s quark suppression was introduced under the requirement that the pertained JETSET parameters mentioned above, must be reduced to the corresponding default values for the $p+p$ collisions at SPS energy, the LUCIAE model with this mechanism works for SPS energy and above only. Therefore we ran LUCIAE without reduction mechanism of s quark suppression for $Au + Au$ collisions at AGS energy but a empirical value of s quark suppression factor ($\text{parj}(2)=0.45$) extracted from experiments [28] was used in stead of default ones in JETSET. The experimental data for $Au + Au$ collisions at AGS energy were taken from [30]. The LUCIAE calculation for $Pb + Pb$ collisions at LHC energy was also given in Tab. 1, in that calculation the $\alpha(\sqrt{s_{nn}})=18$, larger properly than that in $Au + Au$ collisions at RHIC, was assumed. In Tab. 1 the fixed $\alpha(\sqrt{s_{nn}})$ parameter and the obtained fractional variable in string fragmentation function $\langle z \rangle$ (averaged over events), were given as well.

Then we employed the parameter $\alpha(\sqrt{s_{nn}})$ fixed above to calculate the ϕ meson yield, rapidity,

and/or transverse mass distributions in relativistic nucleus-nucleus collisions. The LUCIAE results of ϕ meson yield were compared with the corresponding data in Tab. 2, where the experimental data were taken from [11], [14], and [15] for $Au + Au$ collisions at AGS energy, $Pb + Pb$ at SPS, and $Au + Au$ at $\sqrt{s_{nn}}=130$ GeV, respectively. One sees in Tab. 2 that globally speaking the LUCIAE results are well compatible with experimental data.

In Fig. 1 we compare the LUCIAE results of ϕ meson rapidity (left panel) and transverse mass (right panel, $3.0 < y < 3.8$) distributions with the experimental data in $Pb + Pb$ collisions at SPS energy (data taken from [14]). It is shown in Fig. 1 that the LUCIAE model works somewhat better for the rapidity distribution than the transverse mass distribution in $Pb + Pb$ collisions at SPS energy. The comparison between LUCIAE results and experimental data of ϕ meson transverse mass distributions ($-0.5 < y < 0.5$) in $Au + Au$ collisions at $\sqrt{s_{nn}}=130$ GeV was given in Fig. 2 where data were taken from [15]. One sees in Fig. 1 and 2 that for transverse mass distribution LUCIAE model works better at RHIC energy than SPS since more hard ϕ mesons are produced at RHIC than SPS due to the Schwinger mechanism [31] in particle production of string fragmentation [18].

Fig. 3 (a) gives the fractional variable in string fragmentation function, $\langle z \rangle$, as a function of $\sqrt{s_{nn}}$ from LUCIAE calculations above. A saturation structure at around RHIC energy is observed in Fig. 3 (a). Since the fractional variable z is the fraction of light-cone momentum taken away from the fragmenting string by the produced hadron, it must relate strongly to the nuclear transparency in nucleus-nucleus collisions. The formation time in which string is excited after hard scattering and the hadronic rescattering were considered as dominative factors in baryon stopping (nuclear transparency) [32]. However, one knows that the higher reaction energy the longer formation time, and then the less rescattering and higher nuclear transparency. As for the rescattering itself, the higher reaction energy leads to the more produced particles, and then the more rescattering and lower nuclear transparency. Thus the effects of formation time and the hadronic rescattering on the nuclear transparency cancel each other in certain extent. One might expect that the fractional variable in string fragmentation function is dominant in nuclear transparency and its saturated energy dependence is a qualitative representation of the energy dependence of nuclear transparency.

The corresponding LUCIAE results of charged multiplicity per unit reaction energy $\langle N_{ch} \rangle / \sqrt{s_{AA}}$ in full phase space as a function of $\sqrt{s_{nn}}$ are given in Fig. 3 (b). A trend of saturation seems to be there in Fig. 3 (b) as well. One sees in Fig. 3 (a) that the hadron is

created with higher fractional variable as $\sqrt{s_{nn}}$ is increased and the fractional variable approaches almost to its maximum value one when $\sqrt{s_{nn}}$ gets the LHC energy. On the other hand, Fig. 3 (b) denotes that the efficiency rate of energy to hadron (matter) is decreasing with increase of $\sqrt{s_{nn}}$ and tends also to some kind of saturation. Since the fractional variable equal to one means that only one hadron would be created from fragmentation of an excited string and the number of excited strings (nucleon-nucleon collisions) is limited, the fractional variable approaching to unit must lead to the saturated trend in $\langle N_{ch} \rangle / \sqrt{s_{AA}}$. Those saturation trends might be an indication of the saturation in particle production.

In summary, we employed a hadron and string cascade model, LUCIAE, to study the ϕ meson production in $Au + Au$ and/or $Pb + Pb$ collisions at AGS, SPS, RHIC, and LHC energies. By considering the energy dependence of the model parameter α in the string fragmentation function and adjusting it to the experimental data of charged multiplicity to a certain extent, the model results of ϕ meson yield, rapidity, and/or transverse mass distributions were compatible with the corresponding experimental data for AGS, SPS and RHIC energies. A calculation for the ϕ meson production in $Pb + Pb$ collisions at LHC energy is also given. The obtained fractional variable in string fragmentation function shows a saturation in energy dependence at around RHIC energy. It is discussed that this energy saturation phenomenon could be a qualitative representation of the energy dependence of nuclear transparency.

Finally, the financial supports from NSFC in China (19975075, 10135030, and 10075035) and NRCT in Thailand (1.CH7/2454) are acknowledged.

-
- [1] B. Müller, J. Rafelski, Phys. Rev. Lett. 81, 1066 (1986);
P. Koch, B. Müller, and J. Rafelski, Phys. Rep. 142, 167 (1986).
 - [2] T. Alber, et al., NA35 Collaboration, Z. Phys. C 64, 195 (1994).
 - [3] E. Andersen, et al., NA36 Collaboration, Phys. Lett. B 316, 603 (1993).
 - [4] S. Abatzis, et al., Phys. Lett. B 376, 251 (1996).
 - [5] S. Abatzis, et al., WA94 Collaboration, Phys. Lett. B 400, 239 (1997).
 - [6] I. G. Bearden, et al., NA44 Collaboration, Nucl. Phys. A 610, 175c (1996).
 - [7] M. C. Abreu, et al., NA38 Collaboration, Nucl. Phys. A 638, 483c (1998).
 - [8] E. Andersen et al., WA97 Collaboration, Phys. Lett. B 433, 209 (1998).

- [9] C. Roy, STAR Collaboration, nucl-ex/0111017; hep-ph/0111437; nucl-ex/0201009.
- [10] A. Shor, Phys. Rev. Lett. 54, 1122 (1985).
- [11] B. B. Back, et al., E917 Collaboration, Nucl. Phys. A 661, 506c (1999).
- [12] R. Ferreira, NA38 Collaboration, Nucl. Phys. A 544, 497c (1992).
- [13] M. A. Mazzoni, Nucl. Phys. A 566, 95c (1994).
- [14] S. V. Afanasiev et al., NA49 Collaboration., Phys. Lett. B 491, 59 (2000).
- [15] C. Adler et al., STAR Collaboration, Phys. Rev. C 65, 041901(R) (2002).
- [16] C. M. Ko and B. H. Sa, Phys. Lett. B 258, 6 (1991).
- [17] M. Berenguer, H. Sorge, and W. Greiner, Phys. Lett. B 332, 15 (1994).
- [18] Ben-Hao Sa, Amand Faessler, C. fuchs, An Tai, Xiao-rong Wang, Nu Xu, and E. Zabrodin, Phys. Lett. B 507, 104 (2001).
- [19] S. Pal, C. M. Ko, and Zi-Wei Lin, nucl-th/0202086.
- [20] Sa Ben-Hao and Tai An, Comp. Phys. Comm. 90, 121 (1995); Tai An and Sa Ben-Hao, Comp. Phys. Comm. 116, 353 (1999).
- [21] Tai An and Sa Ben-Hao, Phys. Lett. B 409, 393 (1997).
- [22] Sa Ben-Hao and Tai An, Phys. Rev. C 55, 2010 (1997); Sa Ben-Hao, Wang Xiao-Rong, Tai An, Zhou Dai-Cui, and Cai Xu, Phys. Rev. C 60, 047901 (1999).
- [23] Hong Pi, Z. Phys. C 57, 485 (1993).
- [24] Sa Ben-Hao, Tai An, and Lu Zhong-Dao, Phys. Rev. C 52, 2069 (1995); B. Andersson, An Tai and Ben-Hao Sa, Z. Phys. C 70, 499 (1996).
- [25] B. Andersson and An Tai, Z. Phys. C 71, 155 (1996).
- [26] T. Sjöstrand, Comput. Phys. Commun. 82, 74 (1994).
- [27] W. Ochs, Z. Phys. C 34, 397 (1987).
- [28] J. Cleymans, hep-ph/0201142.
- [29] I. G. Bearden, et al., BRAHMS Collaboration, Phys. Lett. B 523, 227 (2001).
- [30] L. Ahle, et al., E802 Collaboration, Phys. Rev. C 57, 466 (1998).
- [31] J. Schwinger, Phys. Rev. 82, 664 (1951); B. Andersson, G. Gustafson, G. Ingelman, and T. Sjöstrand, Phys. Rep. 97, 31 (1983).
- [32] S. A. Bass, et al., Prog. Part. Nucl. Phys. 41, 255 (1998); nucl-th/9803035.